

# Attenuation of visible solar radiation in the upper water column: A model based on IOPs

ZhongPing Lee<sup>1</sup>, KePing Du<sup>2</sup>, Robert Arnone<sup>1</sup>, SooChin Liew<sup>3</sup>, Bradley Penta<sup>1</sup>

<sup>1</sup>Naval Research Laboratory

Code 7333

Stennis Space Center, MS 39529

[zplee@nrlssc.navy.mil](mailto:zplee@nrlssc.navy.mil)

<sup>2</sup>Research Center for Remote Sensing and GIS

School of Geography

Beijing Normal University

Beijing, 100875, China

<sup>3</sup>Center for Remote Imaging, Sensing and Processing

National University of Singapore

Lower Kent Ridge Road, Singapore 119260

**Abstract** - For many oceanic studies, it is required to know the distribution of visible solar radiation ( $E_{PAR}$ ) in the upper water column. One way to reach this is by remote sensing. This includes two components: First,  $E_{PAR}$  at surface is calculated based on atmosphere properties along with the position of the Sun. Second, the vertical attenuation of  $E_{PAR}$  ( $K_{PAR}$ ) is derived from products of ocean-color remote sensing. Currently,  $K_{PAR}$  is estimated based on chlorophyll concentration ( $[C]$ ) from ocean color. This kind of approach works well for waters where all optical properties can be adequately described by values of  $[C]$ , but will result in large uncertainties for coastal waters where  $[C]$  alone cannot accurately describe the optical properties. In this paper, we present an innovative model that describes  $K_{PAR}$  as a function of water's inherent optical properties (IOP).

## I. INTRODUCTION

Solar radiation in the visible domain ( $E_{PAR}(350 - 700 \text{ nm})$ , measured by downwelling irradiance in this text) encompasses the wavelengths shorter than 700 nm. The pioneer study of *Zaneveld et al.* [1] and subsequent studies [2-4] have demonstrated that the vertical penetration of  $E_{PAR}$  plays an important role in heat transfer of the upper water column.  $E_{PAR}$  at surface can now be adequately estimated from satellite measurements of atmosphere properties. It requires information of water's optical properties to determine the vertical attenuation of  $E_{PAR}$  ( $K_{PAR}$ ) with depth. Historical measurements have shown that  $K_{PAR}$  not only

changes horizontally with constituents in the water [1, 5], but also changes with depth for any water [6, 7].

To represent the steeper than exponential reduction of  $E_{PAR}$  with depth, multiple exponential terms [6, 7] were usually adopted, with an attenuation coefficient (or attenuation depth) assigned for each term. These attenuation coefficients are kept vertically constant, but horizontally vary with *Jerlov* [5] water types. Recently, simple and explicit models have been developed to incorporate satellite-derived chlorophyll concentrations ( $[C]$ ) into the description of the attenuation of  $E_{PAR}$ . When  $[C]$  values are provided via satellite observations of ocean color [8, 9], the partition factors and attenuation coefficients of the terms could be calculated [4].

Such kind of approach works for Case-1 waters - where all optical properties are determined by  $[C]$  alone (with solar zenith angle explicitly or implicitly included) [10, 11]. For non-Case-1 waters, uncertainties arise due to that it is not a constant relationship between  $[C]$  and optical properties. To avoid such limitations associated with  $[C]$ -based models, another approach is to describe the vertical transmittance of  $E_{PAR}$  using water's optical properties [12, 13]. Following this strategy, and because that water's absorption ( $a$ ) and backscattering coefficients ( $b_b$ ) can be adequately derived from ocean-color remote sensing [14-16], we developed a model to describe the vertical transmittance of  $E_{PAR}$  using values of  $a$  and  $b_b$ .

20060623000

## II. Hydrolight SIMULATED $K_{PAR}(z)$

As in earlier studies [17, 18], we used *HydroLight* [19] to get the necessary data sets:  $K_{PAR}(z)$ , and  $a$  and  $b_b$ . Unlike the simulations in *Ohlmann and Siegel* [18] where water's IOPs were determined by [C] only, IOPs in our simulations were simulated with varying [C] and independently varying CDOM and suspended sediments, as described in *Lee et al.* [14] and *IOCCG-OCAG* [20]. Later,  $K_{PAR}(z)$  is modeled as a function of  $a$ ,  $b_b$  and  $z$ . Numerous descriptions can be found regarding simulations by *HydroLight*. [19, 21-23]. The following summarizes the input settings carried out in this study.

The downwelling irradiance at sea surface from the Sun and sky is simulated by the spectral model of *Gregg and Carder* [24].  $a$  and  $b_b$  values at 440 nm varied from 0.02 to 1.9  $m^{-1}$  and 0.002 to 0.115  $m^{-1}$ , respectively, and kept vertically constant. The wavelengths are in a range of 350 – 700 nm with a 10-nm spectral resolution. Five depths (excluding 0 m) were selected for each *HydroLight* run, with depths spread within and beyond the euphotic zone [25]. No bottom reflectance and inelastic scatterings (such as Raman scattering) are included in this study.

## III. MODELING OF $K_{PAR}(z)$

With  $E_{PAR}(z)$  simulated by *HydroLight*,  $K_{PAR}(z)$  is calculated

$$K_{PAR}(z) = \frac{1}{z} \ln \left( \frac{E_{PAR}(0^-)}{E_{PAR}(z)} \right). \quad (1)$$

Figure 1 presents a few examples of  $K_{PAR}(z)$ . Clearly,  $K_{PAR}(z)$  differs significantly for varying water properties. Also, consistent with earlier measurements, subsurface  $K_{PAR}(z)$  changes a lot even for vertically homogeneous waters. This change is due to that water molecules absorb strongly in the longer wavelengths (large absorption coefficients). After photons pass through the subsurface layer (say 3 meters), the absorption is happened in the shorter wavelengths, where absorption coefficients are generally smaller, especially for oceanic waters.

For each vertical variation of  $K_{PAR}(z)$ , it is found that this vertical change could be modeled as,

$$K_{PAR}(IOP, z) = K_1(IOP) + \frac{K_2(IOP)}{(1+z)^{0.5}}. \quad (2)$$

Here  $K_1$  is for the asymptotic value at greater depths, with  $K_2$  more important to the subsurface  $K_{PAR}$  value. IOP here represents different combinations of absorption and backscattering

coefficients. The dotted lines in Figure 1 show Eq.2 modeled  $K_{PAR}(z)$  for those examples. Figure 2 presents the result of *HydroLight*  $K_{PAR}(z)$  versus Eq.2-modeled  $K_{PAR}(z)$ , with the Sun at 30° from zenith. Apparently the modeled  $K_{PAR}(z)$  matches the *HydroLight*  $K_{PAR}(z)$  very well (the average error is 2.2%, with maximum error of is 6.4%). Such results clearly demonstrate that Eq.2 is adequate to describe the vertical change of  $K_{PAR}(z)$ .




Figure 1. Examples of  $K_{PAR}(z)$  for different water properties. The numbers in the box are values of  $a(490)$  (left) and  $b_b(490)$  (right). Symbol represents  $K_{PAR}(z)$  from *HydroLight* simulations, while dotted lines are models from Eq.2.




Figure 2.  $K_{PAR}(z)$  from model (Eq.2) compared with  $K_{PAR}(z)$  from *HydroLight* (30° solar zenith angle), indicating that  $K_{PAR}(z)$  can be well described by Eq.2 with two parameters.

To apply IOP distributions obtained from satellite observation of water color, how  $K_{1,2}$  vary with IOP needs to be known. For the  $K_{PAR}(z)$  data with the Sun at 30° from zenith, it is found that  $K_1$  and  $K_2$  could be well modeled with IOPs at one wavelength *Zaneveld et al.* [12] *Barnard et al.* [13]:

absorption and backscattering coefficients at 490 nm, i.e.,

$$K_1(IOP) = \chi_0 + \chi_1(a(490))^{0.5} + \chi_2 b_b(490), \quad (3a)$$

$$K_2(IOP) = \zeta_0 + \zeta_1 a(490) + \zeta_2 b_b(490). \quad (3b)$$

$\chi_{0,1,2}$  and  $\zeta_{0,1,2}$  are model coefficients.

Since  $K_{PAR}(z)$  also varies with solar altitude, we carried out *HydroLight* simulations with the Sun at  $10^\circ$  and  $60^\circ$  from zenith in order to include solar zenith angle into the model. From these simulations, we got

$$K_1(IOP, \theta_a) = [\chi_0 + \chi_1(a(490))^{0.5} + \chi_2 b_b(490)](1 + \alpha_0 \sin(\theta_a)), \quad (4a)$$

$$K_2(IOP, \theta_a) = [\zeta_0 + \zeta_1 a(490) + \zeta_2 b_b(490)](\alpha_1 + \alpha_2 \cos(\theta_a)). \quad (4b)$$

Here  $\theta_a$  is the solar zenith angle above the surface. Now we got a model that can describe the vertical distribution of  $E_{PAR}$  for different IOPs, depth, and sun angle as

$$T(IOP, z, \theta_a) = \frac{E_{PAR}(z)}{E_{PAR}(0)} = e^{-K_{PAR}(IOP, z, \theta_a)z}. \quad (5)$$

In this model, there are nine model coefficients:  $\chi_{0,1,2}$ ,  $\zeta_{0,1,2}$ , and  $\alpha_{0,1,2}$ . To derive their values,  $T$  values from Eq.5 were fit against  $T$  values from *HydroLight* simulations with the model coefficients derived by least-square curve fitting [4, 18]. Values of derived  $\chi_{0,1,2}$ ,  $\zeta_{0,1,2}$ , and  $\alpha_{0,1,2}$  are provided in Table 1. Figure 3 presents Eq.5 modeled  $T(IOP, z, \theta_a)$  versus  $T(IOP, z, \theta_a)$  determined from *HydroLight* simulations. For those  $T$  values (limiting to the range of  $\sim 0.001$  to  $0.8$ ), bigger errors happened at  $T < 0.003$ , where the effects of  $E_{PAR}$  on heat transfer and photosynthesis in the water column are small. For  $T > 0.003$ , the average error is  $\sim 9\%$ . These results indicate that the simple optical-property-based model (Eq.5) is adequate for describing the vertical profile of  $E_{PAR}(z)$  for different waters.

#### IV. SUMMARY

In this study, an innovative model is developed for describing the vertical transmittance of visible solar radiation ( $E_{PAR}(350 - 700 \text{ nm})$ , measured by downwelling irradiance in this study) in the upper layer of the oceans. Different from the traditional approaches, one exponential term is used for the vertical distribution of  $E_{PAR}$ . Its attenuation coefficient ( $K_{PAR}(z)$ ), however, is modeled as a function of depth instead of vertically constant.  $K_{PAR}(z)$  is also modeled as a function of solar zenith angle and water's optical properties ( $a(490)$

and  $b_b(490)$ ) with data from *HydroLight* simulations. With the availability of  $a(490)$  and  $b_b(490)$  images obtained from satellite remote sensing, this  $K_{PAR}(z)$  model can be adequately incorporated into physical oceanography models to study the effects of visible solar radiation on surface heating [26]. Also, it provides easy and reliable tool to predict the light level at desired depths, needed to plan the  $C^{14}$  incubation for *in situ* measurements of primary production [13].




Figure 3.  $T(z)$  from model (Eq.5) compared with  $T(z)$  from *HydroLight* for three sun angles.

TABLE 1. MODEL COEFFICIENTS FOR  $K_{PAR}(z)$

parameters	values
$\chi_0; \chi_1; \chi_2$	-0.057; 0.482; 4.221
$\zeta_0; \zeta_1; \zeta_2$	0.183; 0.702; -2.567
$\alpha_0; \alpha_1; \alpha_2$	0.090; 1.465; -0.667

#### ACKNOWLEDGEMENT

Support for this study was provided by the Office of Naval Research (P.E. 61153N and N0001405WX20623). The authors thank Dr. Robert F. Chen for assistance in MODTRAN calculation and are grateful to Dr. Curtis Mobley for providing *HydroLight* code and assistance.

#### REFERENCES

1. Zaneveld, J.R.V., J.C. Kitchen, and H. Pak, *The influence of optical water type on the heating rate of a constant depth mixed layer*. J. Geophys. Res., 1981. **86**: p. 6426-6428.
2. Kirk, J.T.O., *Solar heating of water bodies as influenced by their inherent optical properties*. J. Geophys. Res., 1988. **93**(D9): p. 10897-10908.

3. Lewis, M.R., et al., *Influence of Penetrating solar radiation on the heat budget of the equatorial pacific ocean*. *Nature*, 1990. **347**: p. 543-545.
4. Morel, A. and D. Antoine, *Heating rate within the upper ocean in relation to its bio-optical state*. *J. of Physical Oceanography*, 1994. **24**: p. 1652-1665.
5. Jerlov, N.G., *Marine Optics*. 1976, New York: Elsevier.
6. Paulson, C.A. and J.J. Simpson, *Irradiance measurements in the upper ocean*. *J. Phys. Oceanogr.*, 1977. **7**: p. 953-956.
7. Zaneveld, J.R.V. and R.W. Spinrad, *An arc tangent model of irradiance in the sea*. *J. Geophy. Res.*, 1980. **85**(C9): p. 4919-4922.
8. Gordon, H.R. and A. Morel, *Remote assessment of ocean color for interpretation of satellite visible imagery: A review*. 1983, New York: Springer-Verlag. 44.
9. Carder, K.L., et al., *Semianalytic Moderate-Resolution Imaging Spectrometer algorithms for chlorophyll-a and absorption with bio-optical domains based on nitrate-depletion temperatures*. *J. Geophy. Res.*, 1999. **104**: p. 5403-5421.
10. Morel, A., *Optical modeling of the upper ocean in relation to its biogenous matter content (Case I waters)*. *J. Geophy. Res.*, 1988. **93**: p. 10749-10768.
11. IOCCG, *Remote Sensing of Ocean Colour in Coastal, and Other Optically-Complex, Waters*, in *Reports of the International Ocean-Colour Coordinating Group, No.3*, S. Sathyendranath, Editor. 2000, IOCCG: Dartmouth, Canada.
12. Zaneveld, J.R.V., J.C. Kitchen, and J.L. Mueller, *Vertical structure of productivity and its vertical integration as derived from remotely sensed observations*. *Limnol. Oceanogr.*, 1993. **38**: p. 1384-1393.
13. Barnard, A.H., et al., *The determination of PAR levels from absorption coefficient profiles at 490 nm*. *Ciencias Marinas*, 1999. **25**(4): p. 487-507.
14. Lee, Z.P., K.L. Carder, and R. Arnone, *Deriving inherent optical properties from water color: A multi-band quasi-analytical algorithm for optically deep waters*. *Applied Optics*, 2002. **41**: p. 5755-5772.
15. Loisel, H., et al., *Comparison of the ocean inherent optical properties obtained from measurements and inverse modeling*. *Applied Optics*, 2001. **40**: p. 2384-2397.
16. Lyon, P.E., et al., *Chlorophyll biomass in the global oceans: Satellite retrieval using inherent optical properties*. *Applied Optics*, 2004. **43**: p. 5886-5892.
17. Morel, A. and B. Gentili, *Radiation transport within oceanic (case I) water*. *J. Geophy. Res.*, 2004. **109**,C06008, doi:10.1029/2003JC002259.
18. Ohlmann, J.C. and D. Siegel, *Ocean Radiant Heating. Part II: Parameterizing solar radiation transmission through the upper ocean*. *J. of Physical Oceanography*, 2000. **30**: p. 1849-1865.
19. Mobley, C.D., L.K. Sundman, and E. Boss, *Phase function effects on oceanic light fields*. *Applied Optics*, 2002. **41**: p. 1035-1050.
20. IOCCG-OCAG, *Model, parameters, and approaches that used to generate wide range of absorption and backscattering spectra*. 2003, International Ocean Colour Coordinating Group, [http://www.ioccg.org/groups/OCAG\\_data.html](http://www.ioccg.org/groups/OCAG_data.html).
21. Lee, Z.P., et al., *Hyperspectral remote sensing for shallow waters. 1. A semianalytical model*. *Applied Optics*, 1998. **37**: p. 6329-6338.
22. Mobley, C.D., et al., *Comparison of numerical models for computing underwater light fields*. *Applied Optics*, 1993. **32**: p. 7484-7504.
23. Berwald, J., et al., *Influences of absorption and scattering on vertical changes in the average cosine of the underwater light field*. *Limnol. Oceanogr.*, 1995. **40**: p. 1347-1357.
24. Gregg, W.W. and K.L. Carder, *A simple spectral solar irradiance model for cloudless maritime atmospheres*. *Limnol. Oceanogr.*, 1990. **35**: p. 1657-1675.
25. Kirk, J.T.O., *Light & Photosynthesis in Aquatic Ecosystems*. 1994, Cambridge: University Press.
26. Kara, A.B., A.J. Wallcraft, and H.E. Hurlburt, *Sea Surface Temperature sensitivity to water turbidity from simulations of the turbid Black Sea using HYCOM*. *J. Phys. Oceanogr.*, 2005. **35**: p. 33-54.

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p><b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</b></p>						
1. REPORT DATE (DD-MM-YYYY) 07-06-2006		2. REPORT TYPE Conference Proceedings (not refereed)			3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Attenuation of Visible Solar Radiation in the Upper Water Column: A Model Based on IOPs				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER PE0601153N		
				5d. PROJECT NUMBER		
6. AUTHOR(S) ZhongPing Lee, KePing Du, Robert Arnone, SooChin Lioew, Bradley Penta				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER 73-M179-05		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7330-05-5227		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N, Quincy St. Arlington, VA 22217-5660				10. SPONSOR/MONITOR'S ACRONYM(S) ONR		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution is unlimited.						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT For many oceanic studies, it is required to know the distribution of visible solar radiation ( $E_{PAR}$ ) in the upper water column. One way to reach this is by remote sensing. This includes two components: First, $E_{PAR}$ at surface is calculated based on atmosphere properties along with the position of the Sun. Second, the vertical attenuation of $E_{PAR}$ ( $K_{PAR}$ ) is derived from products of ocean-color remote sensing. Currently, $K_{PAR}$ is estimated based on chlorophyll concentration ([C]) from ocean color. This kind of approach works well for waters where all optical properties can be adequately described by values of [C], but will result in large uncertainties for coastal waters where [C] alone cannot accurately describe the optical properties. In this paper, we present an innovative model that describes $K_{PAR}$ as a function of water's inherent optical properties (IOP).						
15. SUBJECT TERMS Visible solar radiation; Ocean-color remote sensing; Remote sensing						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT		18. NUMBER OF PAGES	
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	UL		4	
					19a. NAME OF RESPONSIBLE PERSON Zhongping Lee	
					19b. TELEPHONE NUMBER (Include area code) (228) 688-4873	

## PUBLICATION OR PRESENTATION RELEASE REQUEST

Pubkey: 4494

NRLINST 5600.2

1. REFERENCES AND ENCLOSURES	2. TYPE OF PUBLICATION OR PRESENTATION	3. ADMINISTRATIVE INFORMATION
Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D	( ) Abstract only, published ( ) Book ( ) Conference Proceedings (refereed) ( ) Invited speaker ( ) Journal article (refereed) ( ) Oral Presentation, published ( ) Other, explain	STRN <u>NRL/PP/7330-05-5227</u> Route Sheet No. <u>7330/</u> Job Order No. <u>73-M179-05-5</u> Classification <u>X</u> U <u>  </u> C Sponsor <u>ONR</u> approval obtained <u>X</u> yes <u>  </u> no
Encl: (1) Two copies of subject paper (or abstract)	( ) Abstract only, not published ( ) Book chapter (X) Conference Proceedings (not refereed) ( ) Multimedia report ( ) Journal article (not refereed) ( ) Oral Presentation, not published	

## 4. AUTHOR

Title of Paper or Presentation

Attenuation of Visible Solar Radiation in the Upper Water Column: A Model Based on IOPs

Author(s) Name(s) (First, MI, Last), Code, Affiliation if not NRL

Zhongping Lee, KePing Du, Robert A Arnone, SooChin Liew, Bradley Penta

It is intended to offer this paper to the IGARSS 2005

(Name of Conference)

25-JUL - 29-JUL-2005, Seoul, Korea, Unclassified

(Date, Place and Classification of Conference)

and/or for publication in IGARSS 2005, Unclassified

(Name and Classification of Publication)

(Name of Publisher)

After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a).

It is the opinion of the author that the subject paper (is   ) (is not X) classified, in accordance with reference (b).This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does   ) (does not X) contain any militarily critical technology.This subject paper (has   ) (has never X) been incorporated in an official NRL Report.Zhongping Lee, 7333

Name and Code (Principal Author)

Zhongping Lee  
(Signature)

## 5. ROUTING/APPROVAL

CODE	SIGNATURE	DATE	COMMENTS
Author(s) Lee	<u>Zhongping Lee</u>	<u>4/28</u>	<u>Need by 23 May 05</u>
Section Head Gould	<u>R. H. Gould</u>	<u>4/29/05</u>	<u>Publicly accessible sources used for this publication</u>
Branch Head Robert A Arnone, 7330	<u>Arnone</u>	<u>4/25/05</u>	
Division Head Ruth H. Preller, 7300	<u>R. H. Preller</u>	<u>4/29/05</u>	1. Release of this paper is approved. 2. To the best knowledge of this Division, the subject matter of this paper (has <u>  </u> ) (has never <u>X</u> ) been classified.
Security, Code 7030.1	<u>Ruth H. Preller</u>	<u>5/14/05</u>	1. Paper or abstract was released. 2. A copy is filed in this office.
Office of Counsel, Code 1008.3	<u>W. W. Gould</u>	<u>5/16/05</u>	<u>OSCC-211-5</u> <u>Memo atch</u>
ADOR/Director NCST E.O. Hartwig, 7000	<u>E.O. Hartwig</u>	<u>5/12/05</u>	
Public Affairs (Unclassified/ Unlimited Only), Code 7030.4	<u>Jack Rotunda</u>	<u>5/2/05</u>	
Division, Code			
Author, Code			